



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Black carbon contributes to organic matter in young soils in the Morteratsch proglacial area (Switzerland)

Eckmeier, Eileen ; Mavris, Christian ; Krebs, Rolf ; Pichler, B ; Egli, Markus

Abstract: Most glacier forefields of the European Alps are being progressively exposed since the glaciers reached their maximum expansion in the 1850s. Global warming and climate changes additionally promote the exposure of sediments in previously glaciated areas. In these proglacial areas, initial soils have started to develop so that they may offer a continuous chronosequence from 0 to 150-yr-old soils. The build-up of organic matter is an important factor of soil formation, and not only autochthonous but also distant sources might contribute to its accumulation in young soils and surfaces of glacier forefields. Only little is known about black carbon in soils that develop in glacier forefields, although charred organic matter could be an important component of organic carbon in Alpine soils. The aim of our study was to examine whether black carbon (BC) is present in the initial soils of a proglacial area, and to estimate its relative contribution to soil organic matter. We investigated soil samples from 35 sites distributed over the whole proglacial area of Morteratsch (Upper Engadine, Switzerland), covering a chronosequence from 0 to 150 yr. BC concentrations were determined in fine earth using the benzene polycarboxylic acid (BPCA) marker method. We found that charred organic matter occurred in the whole area, and that it was a main compound of soil organic matter in the youngest soils, where total Corg concentrations were very low. The absolute concentrations of BC in fine earth were generally low but increased in soils that had been exposed for more than 40 yr. Specific initial microbial communities may profit from this additional C source during the first years of soil evolution and potentially promote soil development in its early stage.

DOI: <https://doi.org/10.5194/bg-10-1265-2013>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-86361>

Journal Article

Accepted Version

Originally published at:

Eckmeier, Eileen; Mavris, Christian; Krebs, Rolf; Pichler, B; Egli, Markus (2013). Black carbon contributes to organic matter in young soils in the Morteratsch proglacial area (Switzerland). *Biogeosciences*, 10(3):1265-1274.

DOI: <https://doi.org/10.5194/bg-10-1265-2013>

Black carbon contributes to organic matter in young soils in the Morteratsch proglacial area (Switzerland)

Eileen Eckmeier^{a,*}, Christian Mavris^b, Rolf Krebs^c, Barbara Pichler^b, Markus Egli^b

^aUniversity of Bonn, INRES - Soil Science and Soil Ecology, Nussallee 13, 53115
Bonn, Germany

^bDepartment of Geography, University of Zurich, Winterthurerstrasse 190, 8057
Zurich, Switzerland

^cInstitute of Natural Resource Sciences, Zurich University of Applied Sciences,
Wädenswil, Switzerland

* Corresponding author. Email address: eileen.eckmeier@uni-bonn.de

Abstract

Most glacier forefields of the European Alps are progressively exposed since the glaciers reached their maximum expansion in the 1850s. Global warming and climate changes additionally promote the exposure of sediments in previously glaciated areas. In these proglacial areas, initial soils have started to develop so that they may offer a continuous chronosequence from 0 to 150 year-old soils.

The build-up of organic matter is an important factor of soil formation, and not only autochthonous but also distant sources might contribute to its accumulation in young soils and surfaces of glacier forefields. Only little is known about black carbon in soils that develop in glacier forefields, although charred organic matter could be an important component of organic carbon in Alpine soils.

The aim of our study was to examine whether black carbon is present in the initial

soils of a proglacial area, and to estimate its relative contribution to soil organic matter. We investigated soil samples from 35 sites distributed over the whole proglacial area of Morteratsch, covering a chronosequence from 0 to 150 years. BC concentrations were determined in fine-earth using the benzene polycarboxylic acid (BPCA) marker method. We found that the proportion of BC to total C_{org} was related to the time since the surface was exposed. Soils on surfaces exposed less than 40 years ago contained the highest proportion of BC. The absolute concentrations of BC in fine-earth were generally low but increased in soils that had been exposed for more than 40 years.

Charred organic matter occurred in the whole area, and it was a main compound of soil organic matter in young soils, where total C_{org} concentrations were very low. Specific initial microbial communities consequently may profit from this additional C source during the first years of soil evolution and potentially promote soil development in its early stage.

Keywords: black carbon, BPCA, soil organic matter, glacier forefield, chronosequence, soil development

1. Introduction

Alpine glacier forefields, or proglacial areas, are defined as the area between the present-day glacier front and the terminal moraines deposited in the 1850s, when the glaciers reached their maximum expansion. The most evident developments in Alpine soil formation occur in proglacial areas where new soils form after glaciers retreat and recently formed soils are continuously developing (Egli et al., 2006). Increasing temperatures will lead to a further retreat of glaciers and additional areas will become subject to weathering and formation of soil organic matter. Proglacial

environments are also important for the understanding of global CO₂ cycling on glacial/interglacial timescales as they made up a significant amount of the global land surface during the Quaternary due to the advance and retreat of glaciers and ice sheets (Gibbs and Kump, 1994).

Retreating glaciers often expose barren substrates that become colonised by organisms, beginning the process of primary plant succession which affects the evolution of organic matter in the developing topsoils (Burga et al., 2010). Not only autochthonous but also distant (allochthonous) sources may contribute to the accumulation of soil organic carbon (C_{org}) in young soils and surfaces of glacier forefields. Among these sources are dry and wet atmospheric deposition, faecal deposit, ground-nesting of birds, soot deposit, input due to photo-autotroph organisms, or organic matter of fossil (and preserved) soils (e.g. Bauer et al., 2002; Kim et al., 2005; Arimitsu et al., 2007; Mindl et al., 2007; Xu et al., 2009; Bogdal et al., 2011; Kim et al., 2011; Wientjes et al., 2011). Bernasconi et al. (2011) demonstrated that soil organic matter in initial soils of the Damma glacier forefield contains a relatively high proportion of labile organic compounds which are easily oxidisable and have a short turnover time (annual to decadal). Autochthonous recalcitrant organic matter is not formed in significant amounts in this environment at decadal time scales.

Bardgett et al. (2007) measured changes in the composition of microbial communities and their use of carbon compounds along a 150-year chronosequence in the Austrian Alps. They found that the initial soil microbial communities of the youngest sites were heterotroph organisms that used previously buried recalcitrant carbon, e.g. black carbon, as an energy source. On older sites, after 50 years of exposure, the soil microorganisms respired modern carbon that derived from modern plant growth.

Charred organic matter and soot, or compounds of black carbon (BC), that are released during vegetation fires or fossil fuel burning are ubiquitous components of soils. Several studies demonstrated that BC can be an important compound even in Alpine soils (Bucheli et al., 2004, Eckmeier et al., 2010) or Alpine lake sediments (Bogdal et al., 2011), and in glacial ice cores (Lavanchy et al., 1999; Thevenon et al., 2009), either due to in-situ burning of biomass or via atmospheric deposition. Glaciers can be sources of BC, when they release BC with glacial runoff water that has been incorporated into the ice after melting (Stubbins et al., 2012).

Although the build-up of organic C is an important factor governing the formation of soils and weathering, only very little is known about the presence of BC in very young soils that are developing in glacier forefields. To analyse time trends in such areas, the study of soil chronosequences is an important tool to derive short- to long-term formation rates. The aim of our research was to examine whether BC is an allochthonous source of soil organic matter in the young soils that develop in the proglacial area of the Morteratsch glacier (Upper Engadine, Switzerland) and to estimate its relative contribution (as a function of time) to total organic matter in soils.

2. Materials and methods

2.1 Investigation area

We investigated soils and sediments of the glacier forefield Morteratsch in the Upper Engadine (Switzerland), which is limited by the terminal moraines that have been deposited during the 'Little Ice Age' in the 1850s (Fig. 1). The recent length of this proglacial area is approx. 3 km and it has an area of 1.8 km². The proglacial area is situated in a valley that runs N to S, in an altitude of 1900 m asl to about 2050 m asl. Present climatic conditions are approx. 0.5 °C mean annual temperature and approx. 1000-1300 mm mean annual precipitation (calculated from data from the

meteorological stations Samedan and Bernina). The history of the Morteratsch glacial environment has been studied extensively in geomorphologic and climatic studies (cf. Burga and Perret, 1998; Magny, 1992; Maisch, 1992; Renner, 1982; Gamper, 1985; Fitze, 1982; Patzelt, 1977). The glacial till in the glacier forefield represents the parent material of soil formation. It consists of granitoid and gneissic rock material which underwent a 'green schist' metamorphic event during the high Alpine orogenesis (Büchi, 1994; Spillmann, 1993; Trommsdorff and Dietrich, 1999). Glacial transportation lead to a relatively homogeneous distribution of parent material in the proglacial area.

According to Burga et al. (2010), primary plant succession of the proglacial area started about 7 yr after deglaciation with the *Epilobietum fleischeri* plant community which includes the species *Epilobium fleischeri*, *Oxyria digyna* (only in initial stages), *Linaria alpina*, *Saxifraga aizoides*, and *Rumex scutatus*, and which covered larger areas after about 27 yr. First larch trees, willow and green alder shrubs and the first dwarf-shrubs (e.g. the rust-leaved alpenrose) appeared on areas which have been ice-free for about 12–15 years. Larch-Swiss stone pine stands (*Larici-Pinetum cembrae*) needed more than 150 yr to establish. Only small patches of *Larici-Pinetum cembrae* can be found on the proglacial area while they are dominant in adjacent areas. The succession was not a linear process and influenced by various micro-site dependent factors like soil moisture, grain size, local disturbances or micro-climate that led to a patchy distribution of plant communities.

The dominant soil units (some sites do not have a soil) in the proglacial area are Haplic Fluvisols (Endoskeletal), Skeletic or Lithic Leptosols and Humi-skeletal Leptosols, and Dystric and Endogleyic Cambisols (endoskeletal) (IUSS working group, 2006). The youngest soils showed almost no morphological signs of chemical weathering and alteration products. The development of soils in the Morteratsch area

during 150 years of surface exposure, including soil organic matter formation and mineral weathering processes, have been studied in detail (e.g. Egli et al., 2010; Mavris et al., 2010).

2.2 Methods

Two sets of samples were investigated that had been taken from the whole proglacial area and that cover a chronosequence ranging from 0 to 150 yr (Table 1). Samples S1-10 are ten topsoil samples (soil depth between 1 to max. 12 cm) from ten sites, which are shown in Fig. 1. Close to these sites, material from soil pits has been analysed in detail (Mavris et al., 2010). Samples AS1-29 were sampled from additional sites of the proglacial area at a depth of 0-5 cm.

All samples have been dried at 105°C, sieved (< 2mm) and ball-milled for further analysis.

Total C, nitrogen (N) and hydrogen (H) contents were measured using a C/H/N analyzer (Elementar Vario EL). The organic matter content was determined gravimetrically after dry combustion in a muffle furnace at 550 °C for 6 h (Nocentini et al., 2010). The oxygen (O) content was calculated from the measured data. Total C was considered as organic C (Corg) because CaCO₃ could not be detected in any sample using HCl and only traces (< 0.5%) – if ever – could be observed in DRIFT spectra. The soil is acidic (pH < 6) in the whole proglacial area (Mavris et al. 2010).

The concentration of black carbon (BC) in fine-earth samples was determined as benzene polycarboxylic acids (BPCA) according to the method described by Brodowski et al. (2005). The samples (two replicates) were first treated with trifluoroacetic acid (TFA) to remove polyvalent cations and then digested with HNO₃ at 170 °C for eight hours. The sum of BPCAs in each sample was determined after derivatisation on a gas chromatograph equipped with a flame ionisation detector

(GC-FID). The patterns of benzene rings is dependent on the degree of condensation of the polyaromatic carbon compounds. We used a conversion factor of 2.27 to estimate BC contents from total BPCA-C concentrations. The factor provides a conservative minimum estimate of the true BC contents in soil (Glaser et al., 1998; Brodowski et al. 2005).

Environmental scanning electron microscopy (ESEM) and energy-dispersive spectroscopy (EDS) were performed on three loose granular samples (uncoated) (AS 21, 24, 28) at the Institute for Building Materials (ETH Zurich, Switzerland). The fine-earth fraction of the samples was washed with deionized water and the floating material (density < 1 g/cm³) was collected and air-dried. This enabled the selective extraction of the organic fraction of the sediment, including charcoal material. The analysis was performed using a Dual Beam Quanta 200 3D FEI coupled with EDX, with Dual BSD detector and W emitter operating at an accelerating voltage of 20 kV. The EDS detector is equipped with an ultra-thin window allowing detection of mineral elements and carbon, which provided the elemental composition of the solid phases. Soil colour was measured for dried and homogenised soil samples in triplicates using a spectrophotometer (Konica Minolta CM-5) by detecting the diffused reflected light under standardised observation conditions (2° Standard Observer, Illuminant C). The colour spectra were obtained in the 360 to 740 nm range, in 10 nm increments. The spectral information was converted into the Munsell colour system and the CIELAB Color Space (CIE 1976) using the Software SpectraMagic NX (Konica Minolta). The L* values indicate the extinction of light, or luminance, on a scale from L* 0 (absolute black) to L* 100 (absolute white).

3. Results

3.1 Organic C and nitrogen

The C_{org} concentrations ranged from 1.9-131.0 g kg⁻¹, with an average of 23.6 and a median of 11.5 g kg⁻¹ (Table 2). Figure 2 shows that the C_{org} concentrations in younger soils did not exceed 5 g kg⁻¹ while after about 40 years of surface exposure, the C_{org} concentrations increased and reached values close to 80-100 g kg⁻¹ after about 60 years, and a maximum of 131 g kg⁻¹ at 75 years. There is, however, a strong scatter of C_{org} concentrations that is not only related to the factor time but also to others such as vegetation. The average C_{org} concentrations were higher under the pioneer grass communities (46 g kg⁻¹, 10 sites; vegetation type 2 in Table 1) than under the *Epilobietum fleischeri* sites (17 g kg⁻¹, 15 sites; vegetation type 3). Only three sites were covered by plants belonging to the green alder scrub communities (*Alnetum viridis*), they reached an average of 15 g kg⁻¹.

The N concentrations ranged from 0.2 to 7.0 g kg⁻¹ (mean 1.3, median 0.8 g kg⁻¹). The C/N ratios were between 2 and 58 (mean 19, median 15). For the samples S1-10 that were taken only from topsoil the C/N ratio was between 12 and 36. The distribution of N shows a close relation to the vegetation type, only five sites contain more than 1.4 g N kg⁻¹, of which four are covered by pioneer grass communities (*Geo montani-Nardetum* and *Poion alpinae*, vegetation type 2) and one by green alder scrub communities (*Alnetum viridis*, vegetation type 4).

3.2 Black Carbon

The BC concentrations (calculated with a conversion factor from BPCA-C) varied between 0.2 and 5.6 g kg⁻¹, with an average mean of 1.0 and a median of 0.5 g kg⁻¹ (Table 2). Since the BPCA-C yield exceeded 1.2 g kg⁻¹ C_{org} (without conversion factor), we can exclude competing biogenic sources that could significantly add to the BC concentrations (Brodowski et al., 2005).

The concentration of BC is higher in soils that have been developing since more than

40 years, as is the C_{org} concentration. The BC and C_{org} concentrations were highly correlated ($R^2 = 0.89$, $p < 0.001$). The proportion of BC to total C_{org} was between 23 and 137 g BC $kg^{-1} C_{org}$ (mean 50, median 46 g BC $kg^{-1} C_{org}$). Excluding sample AS6, which was considered an outlier, the relationship with time of surface exposure reaches $R^2 = 0.4$ on a logarithmic scale ($y = -16.4 \ln(x) + 114.35$), as shown in Fig. 3. The youngest soils (< 40 years) contained the highest proportion of BC (68-118 g BC $kg^{-1} C_{org}$), while C_{org} concentrations were low (2-4 g kg^{-1}). The relative distribution of B6CA (6 carboxyls) to total BPCAs was 25% in average, with a tendency to larger amounts of B6CA in older sites (Fig. 4). The patterns of benzene rings were similar for all samples, with the exception of AS6 where the proportion of B6CA reached 50 %.

3.3 Atomic ratios O/C and H/C

The O/C ratios varied from 0.0 to 1.3, the H/C ratios from 0.1 to 2.8. There was no significant relation between O/C ratios and BC concentrations or the proportion of BC at C_{org} . The position of the samples in the van Krevelen plot in Fig. 5 (after Kim et al. 2003) showed that the average atomic ratios for most age groups are typical for plant lignin or cellulose material, with the exception of the youngest age group (0-20 yrs) that was characterized by lower H/C ratios.

3.4 ESEM-EDS

ESEM-EDS allowed the identification of charcoal particles based on their morphology and elemental composition in all analysed samples (AS 21, 24, 28), as shown in Fig. 6. The visual identification was supplemented with the measurement of the atomic ratios of the particles. Following Brodowski et al. (2005), we considered BC as particles that had an O/C ratio of ≤ 0.33 (Stoffyn-Egli et al., 1997) on at least one point

of an observed particle.

3.5 Soil Colour

Soil colour, in particular the Luminance, or brightness, of the soil colour (L^*), varied between 38 and 70 (mean 54, median 55). It was correlated both to C_{org} ($R^2 = 0.545$, $p < 0.001$) and BC concentrations ($R^2 = 0.462$, $p < 0.001$), but not to time.

4. Discussion

Although the environmental conditions in the Morteratsch proglacial area are relatively homogeneous, small scale variability affects soil properties. This is reflected, among others, in the C_{org} concentrations which considerably varied after 30 to 40 years of soil evolution. Small changes such as water content of the substrate, the micro-relief and micro-climate seem to be crucial for both the development of the vegetation and, consequently, also the early evolution of the soils. Two vegetation types persisted over large parts of the chronosequence: pioneer grass communities (Geo montani-Nardetum and *Poion alpinae*, vegetation type 2) and *Epilobietum fleischeri* with single willow shrubs and Alpenrose (vegetation type 3). The C_{org} and N concentrations were affected by these differences in vegetation, the average concentrations were highest under the pioneer grass communities. Oehl et al. (2011) found, for example, that *Epilobium fleischeri* is strongly arbuscular mycorrhizal, but plants in closest distance to the glacier were non-mycorrhizal which finally also influences the C/N ratio in a soil.

Due to the temporal trend, the variability of soil organic matter under all vegetation types was high. The soil colours reflected the heterogeneity of soil formation. Due to their low evolution stage, the soils had a relatively light colour but darkened progressively with increasing C_{org} and BC contents (cf. Eckmeier et al., 2010). The

261 yellow hues showed that the soils were not affected by soil forming processes as
262 brunification yet.

263 BC was present in the whole investigated area, independent from the time since the
264 surface has been exposed, although the concentrations were very low in the
265 youngest soils. The occurrence of BC in remote mountainous environments is not
266 unusual. BC, or, more generally, combustion residues that have been transported as
267 aerosols were deposited on the surfaces of Tibetan glaciers (Xu et al. 2009).
268 Stubbins et al. (2012) found that fossil dissolved organic matter in runoff water of
269 glaciers in Alaska was dominated by aerosols produced during combustion, and that
270 these are a major element in the carbon cycle of glacial environments. BC is also
271 commonly found in the Alpine environments of Switzerland. In the sediments of
272 Lakes Thun, Engstlen and Oberaar, up to one-third of C_{org} was BC (determined by
273 CTO-375; Bogdal et al., 2011). The analysis of ice cores from Colle Gnifetti showed
274 that the deposition of BC increased strongly since the end of the 19th century
275 (Thevenon et al., 2009). The same trend was measured for polycyclic aromatic
276 hydrocarbons (PAHs), which are also produced during burning and which are likely to
277 be absorbed at BC particles. PAH concentrations reached a maximum 1945-1955,
278 and then decreased again. A source assignment using specific PAHs indicated that
279 the ratio of wood and coal burning in contrast to fossil fuel combustion decreased
280 until the 1980s, then the trend reversed (Gabrieli et al., 2010).

281 The source of BC measured in the Morteratsch forefield is charcoal or diagenetic
282 coal. Charcoal particles were found by microscopic inspection, and the BPCA pattern
283 correspond to BPCA pattern of charcoal and coal as described by Roth et al. (2012).
284 The BPCA method, however, would also underestimate the amounts of soot-derived
285 BC (Hammes et al., 2007). Potential sources for the BC in the Morteratsch proglacial
286 area would be wood combustion for heating, charcoal kilns which have been

common in the area until the 20th century, or the railway (Rhaetische Bahn) which is passing in a distance of about 2.5 km from the glacier front that had been equipped with a steam engine until the 1920s.

In two rural areas of Switzerland and in the city of Zurich, burning of wood produced up to one-third of BC emissions (measured during 2.5 years on aerosols using a spectroscopic method) during the winter and 2-10% during summer (Herich et al., 2011). Szidat et al. (2007) even reported contributions of up to 88% of residential wood burning on particulate matter in Alpine valleys during winter. The analysis of 104 soil samples from the Swiss soil monitoring network showed that the BC concentrations had a very uniform distribution because of the uniform deposition of the atmospherically transported BC aerosols (Agarwal & Bucheli, 2010).

Up to now, BC concentrations have not been measured in initial soils of glacier forefields. A small decrease in the proportion of aromatic C with soil age has been found in the Damma glacier forefield using ¹³C-NMR (Dümig et al., 2011), while the proportion of aromatic C in soil organic matter of the Morteratsch glacier forefield, measured using DRIFT, was increasing with time of exposure (Egli et al., 2010). This increase in aromatic compounds was, however, not related to fire-derived organic matter, but rather indicated the presence of condensed and lignin-derived compounds (Poirer et al., 2003).

Charred organic matter was deposited over the whole proglacial area. Local fires could have influenced specific sites, especially S9 where the absolute BC concentration was considerable, or AS6 where the proportion of BC at C_{org} and the proportion of B6CA were highest. The amount of B6CA is slightly but not significantly higher in samples on older surfaces. The youngest soils are characterised by very low C_{org} concentrations, which resulted in a higher proportion of BC in these soils. This is reflected by the H/C ratios which are rather low in the youngest soil samples.

Here, specific initial microbial communities consequently may profit from this additional C source during the first years of soil evolution, as was shown for the Austrian Alps by Bardgett et al. (2007), and potentially promote soil development in its early stage.

5. Conclusion

The developing soils in the Morteratsch proglacial area contained charred organic matter, most likely derived from charcoal or coal, which was distributed over the entire investigated area. It is, however, not known if all BC was deposited by atmospheric deposition, or if the source of BC was material that has accumulated on the glacier and was deposited on the parent material or soil after the ice finally melted. BC concentrations were lowest on surfaces that were exposed during the last 40 years, either due to their lower exposure time or to a reduced input of BC during the last 40 years. BC contributed to total C_{org} which is particularly important at initial soil formation stages, where microorganisms could have used BC as a C source whereas other C_{org} sources provided by vegetation were still scarce or simply lacking.

6. Acknowledgements

This research was supported by the Swiss National Foundation (SNF), project grant numbers 200021-117568 and 200021M_134479, and by the DFG grant EC 401/1-1. We would like to thank I. Wieland, M. Wolf and G. Peschke for assistance with laboratory analyses. We are, furthermore, indebted to Caroline Preston and an unknown reviewer for their helpful comments on an earlier version of the manuscript.

7. References

- 338 Agarwal, T. and Bucheli, T. D.: Is black carbon a better predictor of polycyclic
 339 aromatic hydrocarbon distribution in soils than total organic carbon?, *Environ.*
 340 *Pollut.*, 159, 64–70, 2011.
- 341 Arimitsu, M. L., Piatt, J. F., and Romano, M. D.: Distribution of Ground-Nesting
 342 Marine Birds Along Shorelines in Glacier Bay, Southeastern Alaska: An
 343 Assessment Related to Potential Disturbance by Back-Country Users, Scientific
 344 Investigations Report 2007-5287, UW Dept of the Interior, US Geological Survey,
 345 Reston, Virginia, 2007.
- 346 Bardgett, R. D., Richter, A., Bol, R., Garnett, M. H., Bäumler, R., Xu X., Lopez-
 347 Capel, E., Manning, D. A. C., Hobbs, P. J., Hartley, I. R., and Wanek, W.:
 348 Heterotrophic microbial communities use ancient carbon following glacial retreat,
 349 *Biol. Lett.*, 3, 487–490, 2007.
- 350 Bauer, H., Kasper-Giebl, A., Loflund, M., Giebl, H., Hitenberger, R., Zibuschka, F.,
 351 and Puxbaum, H.: The contribution of bacteria and fungal spores to the organic
 352 carbon content of cloud water, precipitation and aerosols, *Atmos. Res.*, 64, 109–
 353 119, 2002.
- 354 Bernasconi, S. M., Bauder, A., Bourdon, B., Brunner, I., Bünemann, E., Christl, I.,
 355 Derungs, N., Edwards, P., Farinotti, D., Frey, B., Frossard, E., Furrer, G., Gierga,
 356 M., Göransson, H., Gulland, K., Hagedorn, F., Hajdas, I., Hindshaw, R., Ivy-
 357 Ochs, S., Jansa, J., Jonas, T., Kicka, M., Kretzschmar, R., Lemarchand, E.,
 358 Luster, J., Magnusson, J., Mitchell, E. A. D., Olde Venterink, H., Plötze, M.,
 359 Reynolds, B., Smittenberg, R. H., Stähli, M., Tamburini, F., Tipper, E. T., Wacker,
 360 L., Welc, M., Wiederhold, J. G., Zeyer, J., Zimmermann, S., and Zumsteg, A.:
 361 Chemical and biological gradients along the Damma glacier soil chronosequence,
 362 Switzerland, *Vadose Zone J.*, 10, 867–883, 2011.

- 363 Bogdal, C., Bucheli, T. D., Agarwal, T., Anselmetti, F. S., Blum, F., Hungerbühler, K.,
364 Kohler, M., Schmid, P., Scheringer, M., and Sobek, A.: Contrasting temporal
365 trends and relationships of total organic carbon, black carbon, and polycyclic
366 aromatic hydrocarbons in rural low-altitude and remote high-altitude lakes, J.
367 Environ. Monitor., 13, 1316–1326, 2011.
- 368 Brodowski, S., Rodionov, A., Haumaier, L., Glaser, B., and Amelung, W.: Revised
369 black carbon assessment using benzene polycarboxylic acids, Org. Geochem., 36,
370 1299–1310, 2005.
- 371 Bucheli, T. D., Blum, F., Desaulles, A., and Gustafsson, O.: Polycyclic aromatic
372 hydrocarbons, black carbon, and molecular markers in soils of Switzerland,
373 Chemosphere, 56, 1061–1076, 2004.
- 374 Büchi, H.: Der variskische Magmatismus in der östlichen Bernina (Graubünden,
375 Schweiz), Schweiz. Miner. Petrog., 74, 359–371, 1994.
- 376 Burga, C. and Perret, R.: Vegetation und Klima der Schweiz seit dem jüngeren
377 Eiszeitalter, Ott Verlag, Thun, 1998.
- 378 Burga, C. A., Krüsi, B., Egli, M., Wernli, M., Elsener, S., Ziefle, M., and Mavris, C.:
379 Plant succession and soil development on the foreland of the Morteratsch glacier
380 (Pontresina, Switzerland): straight forward or chaotic?, Flora, 205, 561–576, 2010.
- 381 Dümig, A., Smittenberg, R., and Kögel-Knabner, I.: Concurrent evolution of organic
382 and mineral components during initial soil development after retreat of the Damma
383 glacier, Switzerland, Geoderma, 163, 83–94, 2011.
- 384 Eckmeier, E., Egli, M., Hagedorn, F., and Schmidt, M. W. I.: Preservation of fire-
385 derived carbon compounds and sorptive stabilization promote the accumulation of
386 organic matter in black soils of the Southern Alps, Geoderma, 159, 147–155,
387 2010.

- 388 Egli, M., Wernli, M., Kneisel, C., and Haeberli, W.: Melting glaciers and soil
389 development in the proglacial area Morteratsch (Swiss Alps): I. soil type
390 chronosequence, *Arct. Antarct. Alp. Res.* 38, 499–509, 2006.
- 391 Egli, M., Mavris, C., Mirabella, A., Giaccai, D., Kägi, B., and Haeberli, W.: Soil
392 organic matter formation along a chronosequence in the Morteratsch proglacial
393 area (Upper Engadine, Switzerland), *Catena*, 82, 61–69, 2010.
- 394 Fitze, P. F.: Zur Relativdatierung von Moränen aus der Sicht der Bodenentwicklung
395 in den kristallinen Zentralalpen, *Catena*, 9, 265–306, 1982.
- 396 Gabrieli, J., Vallelonga, P., Cozzi, G., Gabrielli, P., Gambaro, A., Sigl, M., Decet, F.,
397 Schwikowski, M., Gäggeler, H., Boutron, C., Cescon, P., and Barbante, C.: Post
398 17th-century changes of European PAH emissions recorded in high-altitude Alpine
399 snow and ice, *Environ. Sci. Technol.*, 44, 3260–3266, 2010.
- 400 Gamper, M.: Morphochronologische Untersuchungen an Solifluktionsszungen,
401 Moränen und Schwemmkegeln in den Schweizer Alpen, *Schriftenreihe Physische*
402 *Geographie*, 17, Geographisches Institut, Universität Zürich, Zürich, 1985.
- 403 Gibbs, M. T. and Kump, L. R.: Global chemical erosion during the last glacial
404 maximum and the present: sensitivity to changes in lithology and hydrology,
405 *Paleoceanography*, 9, 529–543, 1994.
- 406 Glaser, B., Haumaier, L., Guggenberger, G., and Zech, W.: Black carbon in soils: the
407 use of benzenecarboxylic acids as specific markers, *Org. Geochem.*, 29, 811–819,
408 1998.
- 409 Hammes, K., Smernik, R. J., Skjemstad, J. O., Herzog, A., Vogt, U. F., and Schmidt,
410 M. W. I.: Synthesis and characterisation of laboratory-charred grass straw (*Oryza*
411 *sativa*) and chestnut wood (*Castanea sativa*) as reference materials for black
412 carbon quantification, *Org. Geochem.*, 37, 1629–1633, 2006.

- 413 Hammes, K., Schmidt, M. W. I., Smernik, R. J., Currie, L. A., Ball, W. P., Nguyen, T.
 414 H., Louchouart, P., Houel, S., Gustafsson, O., Elmquist, M., Cornelissen, G.,
 415 Skjemstad, J. O., Masiello, C. A., Song, J., Peng, P., Mitra, S., Dunn, J. C.,
 416 Hatcher, P. G., Hockaday, W. C., Smith, D. M., Hartkopf-Froeder, C., Boehmer, A.,
 417 Luer, B., Huebert, B. J., Amelung, W., Brodowski, S., Huang, L., Zhang, W.,
 418 Gschwend, P. M., Flores-Cervantes, D. X., Largeau, C., Rouzaud, J. N., Rumpel,
 419 C., Guggenberger, G., Kaiser, K., Rodionov, A., Gonzalez-Vila, F. J., Gonzalez-
 420 Perez, J. A., La Rosa, J. M. de, Manning, D. A. C., Lopez-Capel, E., and Ding, L.:
 421 Comparison of quantification methods to measure fire-derived (black/elemental)
 422 carbon in soils and sediments using reference materials from soil, water, sediment
 423 and the atmosphere, *Global Biogeochem. Cy.*, 21, GB3016,
 424 doi:10.1029/2006GB002914, 2007.
- 425 Herich, H., Hueglin, C., and Buchmann, B.: A 2.5 year's source apportionment study
 426 of black carbon from wood burning and fossil fuel combustion at urban and rural
 427 sites in Switzerland, *Atmos. Meas. Tech.*, 4, 1409–1420, doi:10.5194/amt-4-1409-
 428 2011, 2011.
- 429 IUSS Working Group: World Reference Base for Soil Resources 2006, 2nd Edn.,
 430 World Soil Resources Reports No. 103, FAO (Food and Agriculture Organisation
 431 of the United Nations), Rome, 2006.
- 432 Kim, J. H., Peterse, F., Willmott, V., Kristensen, D. K., Baas, M., Schouten, S., and
 433 Damste, J. S. S.: Large ancient organic matter contributions to Arctic marine
 434 sediments (Svalbard), *Limnol. Oceanogr.*, 56, 1463–1474, 2011.
- 435 Kim, S., Kramer, R.W., and Hatcher, P. G.: Graphical method for analysis of
 436 ultrahigh-resolution broadband mass spectra of natural organic matter, the Van
 437 Krevelen diagram, *Anal. Chem.*, 75, 5336–5344, 2003.

- 438 Kim, Y., Hatsushika, H., Muskett, R. R., and Yamazaki, K.: Possible effect of boreal
439 wildfire soot on Arctic sea ice and Alaska glaciers, *Atmos. Environ.*, 39, 3513–
440 3520, 2005.
- 441 Lavanchy, V. M. H., Gäggeler, H. W., Schotterer, U., Schwikowski, M., and
442 Baltensperger, U.: Historical record of carbonaceous particle concentrations from
443 a European high-alpine glacier (Colle Gnifetti, Switzerland), *J. Geophys. Res.*,
444 104, 21227–21236, 1999.
- 445 Magny, M.: Holocene lake-level fluctuations in Jura and the northern subalpine
446 ranges, France, regional pattern and climatic implications, *Boreas*, 21, 319–334,
447 1992.
- 448 Maisch, M.: Die Gletscher Graubündens: Rekonstruktion und Auswertung der
449 Gletscher und deren Veränderung seit dem Hochstand von 1850 im Gebiet der
450 östlichen Schweizer Alpen (Bündnerland und angrenzende Regionen),
451 Schriftenreihe Physische Geographie, 32, Universität Zürich-Irchel, Zürich, 1992.
- 452 Mavris, C., Egli, M., Plötze, M., Blum, J., Mirabella, A., Giaccari, D., and Haeberli, W.:
453 Initial stages of weathering and soil formation in the Morteratsch proglacial area
454 (Upper Engadine, Switzerland), *Geoderma*, 155, 359–371, 2010.
- 455 Mindl, B., Anesio, A. M., Meirer, K., Hodson, A. J., Laybourn-Parry, J., Sommaruga,
456 R., and Sattler, B.: Factors influencing bacterial dynamics along a transect from
457 supraglacial runoff to proglacial lakes of a high Arctic glacier, *FEMS Microbiol.*
458 *Ecol.*, 59, 307–317, 2007.
- 459 Nocentini, C., Certini, G., Knicker, H., Francioso, O., and Rumpel, C.: Nature and
460 reactivity of charcoal produced and added to soil during wildfire are particle-size
461 dependent, *Org. Geochem.*, 41, 682–689, 2010.

- 462 Oehl, F., Schneider, D., Sieverding, E. and Burga, C.A.: Succession of arbuscular
463 mycorrhizal communities in the foreland of the retreating Morteratsch glacier in the
464 Central Alps, *Pedobiologia*, 54, 321-331, 2011.
- 465 Patzelt, G.: Der zeitliche Ablauf und das Ausmass postglazialer Klimaschwankungen
466 in den Alpen, in: *Dendrochronologie und postglaziale Klimaschwankungen in*
467 *Europa*, edited by: Frenzel, B., *Erdwissenschaftliche Forschung*, 13, Franz Steiner
468 Verlag, Wiesbaden, 248–259, 1977.
- 469 Poirer, N., Derenne, S., Balesdent, J., Mariotti, A., Massiot, D., and Largeau, C.:
470 Isolation and analysis of the non-hydrolysable fraction of a forest soil and an
471 arable soil (Lacadée, Southwest France), *Eur. J. Soil Sci.*, 54, 243–255, 2003.
- 472 Renner, F.: Beiträge zur Gletschergeschichte des Gotthardgebietes und
473 dendroklimatologische Analysen an fossilen Hölzern, *Schriftenreihe Physische*
474 *Geographie*, 8, Geographisches Institut, Universität Zürich, Zürich, 1982.
- 475 Roth, P. J., Lehndorff, E., Brodowski, S., Bornemann, L., Sanchez-García, L.,
476 Gustafsson, Ö., and Amelung, W.: Differentiation of charcoal, soot and diagenetic
477 carbon in soil: method comparison and perspectives, *Org. Geochem.*, 46, 66–75,
478 2012.
- 479 Spillmann, P.: Die Geologie des penninisch-ostalpinen Grenzbereichs im südlichen
480 Berninagebirge, Ph.D. thesis, ETH Zürich, Switzerland, 1993.
- 481 Stoffyn-Egli, P., Potter, T. M., Leonard, J. D., and Pocklington, R.: The identification
482 of black carbon particles with the analytical scanning electron microscope:
483 methods and initial results, *Sci. Total Environ.*, 198, 211–223, 1997.
- 484 Stubbins, A., Hood, E., Raymond, P. A., Aiken, G. R., Sleighter, R. L., Hernes, P. J.,
485 Butman, D., Hatcher, P. G., Striegl, R. G., Schuster, P., Abdulla, H. A. N.,
486 Vermilyea, A. W., Scott, D. T., and Spencer, R. G. M.: Anthropogenic aerosols as

- 487 a source of ancient dissolved organic matter in glaciers, *Nat. Geosci.*, 5, 198–201,
488 2012.
- 489 Szidat, S., Prévôt, A. S. H., Sandradewi, J., Alfarra, M. R., Sinal, H.-A., Wacker, L.,
490 and Baltensperger, U.: Dominant impact of residential wood burning on particulate
491 matter in Alpine valleys during winter, *Geophys. Res. Lett.*, 34, L05820,
492 doi:10.1029/2006GL028325, 2007.
- 493 Thevenon, F., Anselmetti, F. S., Bernasconi, S. M., and Schwikowski, M.: Mineral
494 dust and elemental black carbon records from an Alpine ice core (Colle Gnifetti
495 glacier) over the last millennium, *J. Geophys. Res.*, 114, D17102,
496 doi:10.1029/2008JD011490, 2009.
- 497 Trommsdorff, V. and Dietrich, V.: *Grundzüge der Erdwissenschaften*, 6th Edn., vdf-
498 Verlag, Zurich, Switzerland, 1999.
- 499 Wientjes, I. G. M., Van de Wal, R. S. W., Reichert, G. J., Sluijs, A., and Oerlemans,
500 J.: Dust from the dark region in the western ablation zone of the Greenland ice
501 sheet, *The Cryosphere*, 5, 589–601, doi:10.5194/tc-5-589-2011, 2011.
- 502 Xu, B., Cao, J., Hansen, J., Yao, T., Joswila, D. R., Wang, N., Wu, G., Wang, M.,
503 Zhao, H., Yang, W., Liu, X., and He, J.: Black soot and the survival of Tibetan
504 glaciers, *P. Natl. Acad. Sci. USA*, 106, 22114–22118, 2009.

505

Tables

Table 1. General properties of the investigated sites in the proglacial area.

Site	Year of exposure	Exposure time	Horizon	Depth	Skeleton	Soil type (WRB)	Vegetation ^a
		yr		cm	wt-%		
S1	1870	140	A	0-6	41	Humi-skeletal Leptosol	4
AS1	1870	140	A	0-5	n.d.	Humi-skeletal Leptosol	2
S2	1880	130	A	0-10	64	Humi-skeletal Leptosol	4
AS2	1880	130	A	0-5	n.d.	Humi-skeletal Leptosol	2
AS3	1890	120	A	0-5	n.d.	Humi-skeletal Leptosol	5
AS4	1890	120	A	0-5	n.d.	Skeletal Leptosol	2
S3	1900	110	A	0-5	54	Humi-skeletal Leptosol	3
AS5	1900	110	A	0-5	n.d.	Skeletal Leptosol	7
S4	1910	100	A	0-1	55	Humi-skeletal Leptosol	3
AS6	1910	100	AC	0-5	n.d.	Skeletal Leptosol	2
AS8	1920	90	AC	0-5	n.d.	Skeletal Leptosol	6
S10	1930	80	A1	0-2	49	Humi-skeletal Leptosol	3
AS9	1930	80	AC	0-5	n.d.	Skeletal Leptosol	4
S9	1935	75	O	0-3	44	Humi-skeletal Leptosol	2
S5	1940	70	A1	0-1	7	Humi-skeletal Leptosol	2
AS10	1940	70	A	0-5	n.d.	Humi-skeletal Leptosol	2
AS11	1945	65	A	0-5	n.d.	Humi-skeletal Leptosol	3
AS12	1945	65	A	0-5	n.d.	Skeletal Leptosol	3
S8	1950	60	O/A	0-12	63	Skeletal Leptosol	2
AS13	1950	60	A	0-5	n.d.	Skeletal Leptosol	3
AS14	1950	60	A	0-5	n.d.	Humi-skeletal Leptosol	3
S6	1960	50	A	0-3	64	Skeletal Leptosol	3
S7	1960	50	A	0-4	26	Skeletal Leptosol	2
AS15	1965	45	A	0-5	n.d.	Skeletal Leptosol	2
AS16	1965	45	(A)C	0-5	n.d.	Skeletal Leptosol	3
AS17	1970	40	AC	0-5	n.d.	Skeletal Leptosol	3
AS18	1970	40	AC	0-5	n.d.	Skeletal Leptosol	3
AS19	1975	35	(A)C	0-5	n.d.	-	3
AS20	1975	35	(A)C	0-5	n.d.	-	3
AS21	1975	35	(A)C	0-5	n.d.	-	3
AS22	1980	30	(A)C	0-5	n.d.	-	0
AS23	1980	30	(A)C	0-5	n.d.	-	3
AS24	1990	20	(A)C	0-5	n.d.	-	0
AS28	2007	3	C	0-5	n.d.	-	0
AS29	2007	3	C	0-5	n.d.	-	0

n.d. = not determined

^a 0 = no vegetation; 2 = Pioneer grass communities (*Geo montani-Nardetum* and *Poion alpinae*); 3 = *Epilobietum fleischeri* with single willow shrubs and Alpenrose; 4 = Green alder scrub (*Alnetum viridis*) (*Alnus viridis* and tall perennial herbs, *Salix spec.*, *Poa spec.*, *Deschampsia caespitosa*, *Avenella flexuosa*, *Nardus stricta*, *Festuca spec.*, *Phleum rhaeticum*, *Anthoxanthum alpinum*, *Calamagrostis villosa*); 5 = Grass heath on moister soils (grass species e.g. *Festuca violacea*, *Calamagrostis villosa*, *Phleum rhaeticum*, *Poa alpina*); 6 = Boulder plant communities, partially *Epilobietum fleischeri* (*Epilobium fleischeri*, *Adenostyles leucophylla*, *Rumex scutatus*, *Dryopteris spec.*, *Athyrium spec.*, *Gymnocarpium dryopteris*, *Polystichum lonchitis*); 7 = Rock vegetation (e.g. *Agrostis rupestris*, *Silene rupestris*, *Sempervivum arachnoideum*)

519 Table 2. Spectral and chemical properties of the fine-earth fraction (< 2mm).

Site	Exposure time	Lumi- nance	Munsell colour			C _{org}	N	C/N	O/C	H/C	BC ^a	BC ^a	B6CA
	yr	L *	Hue	Value	Chroma	g kg ⁻¹	g kg ⁻¹	Weight ratio	Atomic ratio		g C kg ⁻¹ C _{org}	g kg ⁻¹	%
S1	140	57.9	2.5 Y	5.6	1.5	7.1	0.6	12	0.7	1.0	49.7	0.4	27.8
AS1	140	47.5	1.6 Y	4.6	1.5	61.2	4.6	13	0.6	1.8	43.1	2.5	26.3
S2	130	40.2	0.8 Y	3.9	1.1	64.1	3.0	22	0.5	0.8	43.2	2.8	22.8
AS2	130	48.1	2.2 Y	4.7	1.8	18.1	1.2	15	0.5	1.6	48.8	0.8	19.5
AS3	120	58.3	2.9 Y	5.7	1.5	7.3	0.3	23	0.7	0.3	54.5	0.4	43.9
AS4	120	54.8	2.3 Y	5.3	1.7	9.4	0.4	24	0.8	0.4	49.9	0.5	38.5
S3	110	51.6	2.3 Y	5.0	1.4	30.3	1.0	29	0.4	1.3	33.1	1.0	27.7
AS5	110	47.2	1.9 Y	4.6	1.7	14.2	1.4	10	1.0	0.7	58.1	0.8	28.4
S4	100	39.0	1.4 Y	3.8	1.5	54.5	1.2	47	0.7	1.7	25.8	1.4	20.7
AS6	100	62.2	3.1 Y	6.1	1.3	3.2	0.7	4	1.3	2.2	136.7	0.4	49.9
AS8	90	63.9	2.5 Y	6.2	1.6	3.9	0.4	10	0.2	1.7	n.d.	n.d.	n.d.
S10	80	38.4	1.4 Y	3.7	1.3	47.9	1.3	36	0.9	1.6	44.9	2.1	14.5
AS9	80	62.6	3.4 Y	6.1	1.4	5.3	0.6	9	1.2	1.5	44.7	0.2	28.8
S9	75	39.4	0.5 Y	3.8	1.4	131.0	7.0	19	0.7	1.6	42.8	5.6	13.5
S5	70	57.1	2.5 Y	5.6	1.1	31.7	0.8	42	0.6	1.1	40.2	1.3	26.1
AS10	70	49.5	2.0 Y	4.8	1.8	29.9	1.1	28	0.5	0.8	61.4	1.8	23.6
AS11	65	49.4	2.1 Y	4.8	1.7	11.6	0.8	14	0.7	0.5	46.6	0.5	24.4
AS12	65	43.8	1.8 Y	4.3	1.6	12.6	0.7	19	0.7	0.4	30.2	0.4	24.1
S8	60	45.1	1.7 Y	4.4	1.2	89.7	3.8	24	0.6	1.5	25.9	2.3	14.8
AS13	60	47.6	2.2 Y	4.6	1.3	11.5	0.4	30	0.5	0.9	27.1	0.3	25.3
AS14	60	58.5	2.7 Y	5.7	1.4	18.1	0.3	58	0.6	1.4	23.7	0.4	22.0
S6	50	48.1	1.4 Y	4.7	1.8	37.0	1.1	34	0.3	1.6	27.9	1.0	33.6
S7	50	43.2	1.6 Y	4.2	1.7	64.5	4.3	15	0.6	1.6	23.0	1.5	21.6
AS15	45	49.7	2.3 Y	4.8	1.6	18.3	0.3	57	0.8	1.6	48.0	0.9	16.6
AS16	45	61.7	2.7 Y	6.0	1.2	4.9	1.4	3	1.2	1.4	46.4	0.2	24.3
AS17	40	58.6	2.2 Y	5.7	1.6	6.0	1.0	6	0.3	1.2	50.1	0.3	23.2
AS18	40	59.1	2.6 Y	5.8	1.5	9.3	0.4	25	0.8	0.9	35.6	0.3	20.4
AS19	35	69.6	2.4 Y	6.8	1.4	2.3	1.0	2	0.2	1.4	n.d.	n.d.	n.d.
AS20	35	59.9	3.0 Y	5.8	1.5	4.1	1.3	3	0.0	1.8	67.7	0.3	22.3
AS21	35	54.7	3.1 Y	5.3	1.5	4.0	0.7	5	0.0	0.1	n.d.	n.d.	n.d.
AS22	30	59.7	3.7 Y	5.8	1.4	2.9	0.3	10	0.3	2.1	69.2	0.2	22.7
AS23	30	64.7	4.2 Y	6.3	1.3	2.6	0.2	11	0.6	0.3	73.7	0.2	22.9
AS24	20	63.8	3.8 Y	6.2	1.4	2.9	0.4	7	0.0	2.2	n.d.	n.d.	n.d.
AS28	3	62.5	4.3 Y	6.1	1.3	1.9	0.7	3	0.9	0.4	117.9	0.2	22.6
AS29	3	60.3	4.5 Y	5.9	1.2	2.1	0.3	8	0.7	2.8	n.d.	n.d.	n.d.

^a Black carbon was calculated by multiplying BPCA-C data with a conversion factor of 2.27 (Brodowski et al. 2005).

520
521
522
523

Figure Captions






Fig. 1: Location of the Morteratsch glacier forefield with isochrones of glacier retreat, major soil units and position of the topsoil sampling sites, which were analysed in detail regarding soil chemistry and mineralogy (S1–10; Mavris et al., 2010).



529

530 Fig. 2: C_{org} concentrations (in g kg⁻¹) in all samples as a function of surface exposure
 531 time and vegetation cover. The vegetation types are related to the numbers given in
 532 Table 1.

533



534

535 Fig. 3: Relative proportion of BC in total organic carbon as a function of time for all

536 samples. Sample AS6 was classified as an outlier and removed from the regression.

537




Fig. 4: Average patterns of benzene rings in proportion to total BPCA-C, showing a trend to larger contributions of B6CA in soil material on older surfaces. Outlier AS6 is included in the age group 100-120. The standard error is given for samples of the corresponding group of time of exposure.




Fig. 5: Van Krevelen diagram showing the H/C and O/C atomic ratios of soils (mean values \pm SD of age classes). Grey shaded areas indicate areas of organic compounds after Kim et al. (2003) and Hammes et al. (2006).




Fig. 6: Environmental scanning electron microscopy (ESEM) and energy-dispersive spectroscopy (EDS) on organic particles (charcoal) with a density $< 1 \text{ g cm}^{-3}$. A) exposure age three years (AS29); B) exposure age 20 years (AS24). BSE = back-scattered electrons.